



Article Side

Improving Breast Symmetry through Breast Enlargement by [Terry Bayer](#)

Article published on March 20th 2012 | [Health](#)

Women have different breast measurements and most if not all females have asymmetrical breasts or breasts that are not of the same size. Most ladies who have lopsided breasts are those who experienced breastfeeding babies. Breastfeeding women usually offer the same breast during feeding time resulting in the uneven size. But, this must not be a cause for worry because there are various procedures that can help reduce breast asymmetry. These techniques are simple and easily tolerable for a lot of patients. Some doctors recommend that women wait until the age of 20, when their breasts reached their full development, before they undergo any breast asymmetry procedure or any cosmetic procedure for the breast.

Some ladies have well-developed breasts but are not in proportion with their bodies. Others have smaller breasts but would choose to have them bigger so they can wear clothes that appear better with fuller breasts. Developments in cosmetic surgery now make it possible for full-breasted women to have breasts that are improved; and for small-breasted women to get fuller ones. The treatments are both quick and harmless.

For women with full but uneven or lopsided breasts, a technique to reduce breast asymmetry can be done. This may also involve decrease in nipple size. The reconstructed breasts are lessened and lifted to look well-shaped.

Women with smaller breasts can opt for implants to correct breast asymmetry. Implants can either be silicone or saline breast implants. Silicone implants are filled with silicone gel that appears like human fat. Some women prefer this type of implant over the other as it feels and looks more comfortable. Saline implants are filled with saline solution during the time of operation. Other patients say saline implants are not as soft and look unnatural. Patients can seek advice from their doctors on which is the less dangerous and better selection for them.

People who go through either treatment must put on special bras to help decrease breast swelling. These bras will give the breasts ample support since inflammation and tenderness are expected for at least two days after the surgical treatment. Doctors recommend clients who have undergone breast augmentation Los Angeles clinics offer to wear the special bra for a couple of weeks and keep from doing strenuous activities like lifting or pushing.

Surgeons in Los Angeles are known to have more expertise in correcting breast asymmetry because of the wide clientele they have. They can also conduct breast implant revision procedures if clients report that they are not fully satisfied with the breast procedure they got.

Article Source:

<http://www.articleside.com/health-articles/improving-breast-symmetry-through-breast-enlargement.htm> - [Article Side](#)

[Terry Bayer](#) - About Author:

For more details, search a [breast asymmetry](#), a [breast augmentation Los Angeles](#), and a [breast implant revision](#) in Google for related information.

Article Keywords:

breast asymmetry, breast augmentation Los Angeles, breast implant revision

You can find more [free articles](#) on [Article Side](#). Sign up today and share your knowledge to the community! It is completely FREE!