



Article published on March 20th 2012 | [Health](#)

Ladies have different breast measurements and many if not all females have asymmetrical breasts or breasts that are not of the same specifications. Most females who have uneven breasts are those who experienced breastfeeding. Breastfeeding women usually give the same breast during feeding time causing uneven size. But, this should not be a cause for worry as there are numerous procedures that can help correct breast asymmetry. These methods are simple and easily tolerable for many patients. Some doctors advise that women wait until age 20, when their breasts have fully developed, before they undergo any breast asymmetry treatment or any cosmetic procedure for the breast.

Some women have developed breasts but are not even. Others have small breasts but would rather have them bigger to allow them to wear clothes that appear better with fuller breasts. Advancements in plastic surgery now make it possible for full -breasted women to get breasts that are well-shaped; and for small-breasted women to have fuller ones. The treatments are both quick and safe.

For women with developed but uneven or lopsided breasts, a technique to eliminate breast asymmetry can be made. This is may also involve decrease in nipple size. The reconstructed breasts are lessened and lifted to look well-shaped.

Females with smaller breasts can choose implants to correct breast asymmetry. These implants can be silicone or saline implants. Silicone implants are packed with silicone gel that appears like human fat. Some ladies prefer this type of implant over the other as it feels and looks more natural. Saline implants are filled with saline solution during operation. Other patients say saline implants are not as soft and appear unnatural. Patients can talk to their doctors on which is the less dangerous and better option for them.

Patients who go through any of the implant options must wear special bras to help lessen breast swelling. These bras will give the breasts sufficient support since inflammation and pain are expected not less than two days after the treatment. Doctors advise clients who have undergone breast augmentation Los Angeles clinics offer to put on the special bra for a number of weeks and refrain from doing physically demanding activities like lifting or pushing.

Surgeons in Los Angeles are known to have more expertise in correcting breast asymmetry because of the wide clientele they have. They are also able to carry out breast implant revision techniques in case clients declare that they are not fully satisfied with the breast procedure they experienced.

Article Source:

<http://www.articleside.com/health-articles/reconstructing-breasts-via-breast-augmentation-solutions.htm> - [Article Side](#)

[Terry Bayer](#) - About Author:

For more details, search a [breast asymmetry](#), a [breast augmentation Los Angeles](#), and a [breast implant revision](#) in Google for related information.

Article Keywords:

breast asymmetry, breast augmentation Los Angeles, breast implant revision

You can find more [free articles](#) on [Article Side](#). Sign up today and share your knowledge to the community! It is completely FREE!